

Acacia ligulata **Umbrella Wattle (Prostrate)**









Females (top) and male Twospotted Line-Blue butterflies

(Fagg, M, Australian Plant Image Index)

(Murfet, D, Seeds of South Australia)

(SA Butterflies & Moths)

Form: Ground cover **Height:** up to 0.2 m **Spread:** up to 2 m

Flowers: Yellow

Type: Globular

Flowering time: Spring



Seed collecting: collect mature pods that are

Seed treatment: place seed into a container

and pour on water that has just been boiled.

Soak overnight. Remove and sow immediately.

turning brown, with hard, dark seeds inside.

Soils: sandy, clay or loam; acidic, alkaline or neutral

Sowing requirements: cover seed to its own depth with damp soil and firm gently. Then cover with a layer of gravel. Sow in late Spring-

Position: full sun

Supplementary watering: minimal

Germination rate: 4 to 6 weeks

Uses: A fast growing screen and soil stabiliser. The roots are habitat for witchetty grubs. Indigenous people used the seeds to make flour. Food host to Two-spotted Line-blue (Nacaduba biocellata) caterpillars.

Notes: The shallow roots can inhibit growth to plants around it.

Sources: Botanical Gardens of South Australia, Seeds of South Australia, TreeProject, Trees For Life Tree Scheme: Grower's Handbook; SA Butterflies & Moths