

Dianella revoluta Black Anther Flax Lily







Adult Southern Grass-dart

(Fisher, R, SA Butterflies & Moths)



Adult Greenish Grass-dart
(Grund, R, SA Butterflies & Moths)

(Elder, B, iNaturalist Australia)

Form: Sedge or Flax **Height:** 0.2 – 0.3 m

(Best, R, iNaturalist Australia)

Spread: 0.4 - 0.5 m **Flowers:** Blue

Type: Star

Flowering time: Spring, Summer

Germination rate: 3 to 8 weeks

Ecological functions:







Soils: sand, clay or loam; acidic, alkaline or neutral

Seed collecting: wearing gloves, collect fruits that are losing colour or drying off. Allow to dry. Once dry rub firmly to dislodge shiny black seeds. Store in cool dry area.

Seed treatment: seed packet will contain either pure seed or seed still in the fruit. Split fruits to release seed if necessary. Stratification by cold treatmnet can help break dormancy.

Sowing requirements: cover with 2 mm of damp soil and firm gently. Then cover with a layer of gravel. Sow in late Summer to early <u>Spring. Requires</u> well-drained soils.

Position: part shade, full sun.

Supplementary watering: minimal

Uses: border plant for median strips and roundabouts. Can be used for erosion control in exposed coastal locations. Rare food host to Southern Grass-dart (*Ocybadistes walkeri hypochlorus*) and Greenish Grass-dart (*Ocybadistes walkeri sothis*) caterpillars.

Notes: unedible, harmful if eaten. Do not

Sources: Botanical Gardens of South Australia; Seeds of South Australia; Trees For Life Tree Scheme: Grower's Handbook'; SA Butterflies & Moths; "What Seed is that?"