

Xanthorrhoea semiplana

Grass Tree









Seeds attract
Yellow-tailed
Black-Cockatoos
(pictured), and
nectar attracts
other birds.
(Lawrenz, R,
iNaturalist
Australia)

(Hura, M, iNaturalist Australia)

(Murfet, D, Seeds of South Australia)

(Murfet, D, Seeds of South Australia)

Form: Medium shrub

Height: 1-3 m **Spread:** 1-2 m

Flowers: Cream Type: Spikes

Flowering time: Spring, Winter

Germination rate: 4 to 52 weeks

Ecological functions:



Seed collecting: place material underneath the

the flowering stalk and bend it over. Shake and

many mature, black seeds will fall out. Near

maturing fruits can be cut off and placed in warm area to fully dry. Ample seeds will fall out.

Pass material through screens to clean fully,

add insecticide dust before storing away.





Soils: sand, clay or loam; acidic, alkaline or neutral

Sowing requirements: sprinkle seeds over propagating tray. Cover lightly to approximately 4 mm with soil or fine gravel. Keep moist and place in open, sunny area. Sow from early Spring or in Autumn. Requires good

Position: full sun

drainage.

Supplementary watering: minimal

Uses: ornamental. Adapted to fires, trunk blackens but does not get damaged. Very slow growing. European settlers harvested the resinous gum to make varnishes and lacquers. Susceptible to and a good indicator of *Phytophthora cinnamomi*. Notes: written approval required prior to planting in streets or roads. Do not plant closer than 2 m to any sewer main or connection.

Seed treatment: none